

<b>Cardiopulmonary</b>	<b>Musculoskeletal</b>	<b>Skin</b>	<b>Gastrointestinal</b>	<b>Elimination</b>
<ul style="list-style-type: none"> <li>• Ventilator dependencies</li> <li>• Tracheostomy</li> <li>• Weakened cough &amp; gag reflex</li> <li>• Difficulty Swallowing</li> <li>• Narrowing of the trachea</li> <li>• Recurrent regurgitation</li> <li>• Increase risk of aspiration</li> <li>• Pneumonia</li> <li>• Respiratory infections</li> </ul> <p>Abnormal heart rates</p> <ul style="list-style-type: none"> <li>• Abnormal rhythm</li> <li>• Hypertension</li> </ul>	<ul style="list-style-type: none"> <li>• Peripheral nerve injuries</li> <li>• Spasticity</li> <li>• Contractures</li> <li>• Heterotopic ossification</li> <li>• Deep venous thrombosis</li> <li>• Reflex sympathetic dystrophy</li> <li>• Phlebitis</li> <li>• Cellulitis</li> </ul>	<ul style="list-style-type: none"> <li>• Lacerations</li> <li>• Abrasions</li> <li>• Acne</li> <li>• Profuse sweating</li> <li>• Medication reactions</li> <li>• Worsened skin conditions</li> <li>• Pressure ulcers</li> </ul>	<ul style="list-style-type: none"> <li>• Change in nutritional needs</li> <li>• Increased calorie needs</li> <li>• Swallowing disorders</li> <li>• Gastronomy tubes</li> <li>• Impaired cognition from gastrointestinal medications</li> </ul>	<ul style="list-style-type: none"> <li>• Unaware of the need to eliminate</li> <li>• Constipation or diarrhea</li> <li>• Loss of control over elimination</li> <li>• Disinhibited neurogenic bladder</li> <li>• Need for catheter</li> </ul>